

JOURNAL OF DIFFERENTIAL EQUATIONS 10, 262-269 (1971)

On Periodic Solutions of the System

$$\dot{x} = y - F(x), \quad \dot{y} = -g(x)^*$$

PETER J. PONZO

Department of Applied Mathematics, University of Waterloo, Waterloo, Ontario

AND

NELSON WAX

Department of Electrical Engineering, University of Illinois, Urbana, Illinois 61801

Received June 23, 1970

1. INTRODUCTION

The equation

$$\ddot{y} + \mu \sin \dot{y} + y = 0, \quad \left(\cdot \equiv \frac{d}{dt} \right) \quad (1)$$

or its equivalent representation

$$\dot{x} = y - \mu \sin x, \quad \dot{y} = -x \quad (2)$$

furnishes an approximate description of the behavior of a second-order phase-locked loop, an electrical circuit used extensively in a variety of applications, notably in space communication.

The generalization of (2)

$$\begin{aligned} \dot{x} &= y - F(x), \\ \dot{y} &= -g(x), \end{aligned} \quad (3)$$

yields descriptions of the motion of Froude's pendulum, of the angular oscillations of a synchronous motor, and of the variation in electrical phase in a power distribution system, when F or g or both are taken to be oscillatory.

* This research was supported in part by a grant from the National Research Council Ottawa, Canada, and in part by the Joint Services Electronics Program, Contract DAAB-07-67-C-0199.

Despite the interest in (2) and (3) occasioned by these applications, it is only recently that Hochstadt and Stephan [1] first proved the existence of an infinite number of limit cycles for (2) when $|\mu|$ is sufficiently small. D'heedene [2] then showed that (2) possesses an infinite number of periodic solutions for all real μ . Comstock [3] generalized D'heedene's results, replacing $\sin x$ by an odd oscillatory $F(x)$ which satisfies certain hypotheses, given in Section 2.

Here, we extend this work to system (3) without including, however, oscillatory $g(x)$. We prove, in Section 2, the existence of a denumerable set of periodic orbits when $F(x)$ satisfies Comstock's hypotheses and $g(x)$ is odd and suitably restricted. We turn, in Section 3, to even $F(x)$, odd $g(x)$, and show that under some relatively mild hypotheses, system (3) will possess a nondenumerable set of periodic solutions.

2. PERIODIC SOLUTIONS: $F(x)$ AND $g(x)$ ODD

Let

H_1 :

(a) $F(x) = -F(-x)$; $F(x) \in C^2(-\infty, \infty)$; with $F'(x) = f(x)$, $|F(x)| \leq A$, and $|f(x)| \leq B$ for all x , with $yF(y) > 0$ for y sufficiently small.

(b) $F(x)$ be oscillatory; with $\{z_k\}^\infty$ the sequence of positive zeros of $F(x)$ one requires $\int_{z_n}^{z_{n+1}} |F(x)| dx$ to be a nondecreasing function of n . (These are Comstock's [3] hypotheses concerning $F(x)$.)

Furthermore, assume

H_2 :

(a) $g(x) = -g(-x)$; $g(x) \in C^1(-\infty, \infty)$, and, for some a , $g'(x) \geq a > 0$ for all x .

(b) with $G(x) = \int_0^x g(u) du$, that there exists an $x^* \geq A/\sqrt{a}$ such that $G(x)/g(x) \geq 2A\{B + \sqrt{a}\}/a$ for all $x \geq x^*$.

Consider the simple closed convex curves defined by

$$\lambda(x, y, C) = \frac{1}{2}(y - C)^2 + G(x) = \text{const.}$$

The time rate of change of λ along a solution trajectory is given by $\dot{\lambda} = g(x)[C - F(x)]$. If $C = A$, then, since $|F(x)| \leq A$, solutions in $x \geq 0$ cross these ovals from their interiors to their exteriors. Similarly, the ovals $\lambda(x, y, -A) = \text{constant}$ are crossed toward their interiors, in $x \geq 0$.




FIG. 1. The upper branch y_U and the lower branch y_L of a trajectory, together with two bounding ovals, are shown for $x \geq 0$.

We use these contours (which are analogous to the circles of D'heedene) to obtain the following bounds on the upper branch y_U and the lower branch y_L of a solution trajectory (see Fig. 1).

$$\begin{aligned} -A + \{2[G(z) - G(x)]\}^{1/2} &\leq y_U \leq A + \{2[G(z) - G(x)]\}^{1/2}, \\ -A - \{2[G(z) - G(x)]\}^{1/2} &\leq y_L \leq A - \{2[G(z) - G(x)]\}^{1/2}. \end{aligned} \quad (4)$$

Consider now the change in $\lambda(x, y, 0)$ as a solution proceeds from P to Q (Fig. 1). One has $(d/dt) \lambda(x, y, 0) = -g(x)F(x)$, so that $d\lambda = -(gF/y - F) dx$. Thus

$$\lambda(R) - \lambda(P) = \int_0^z -\frac{gF}{y_U - F} dx, \quad \text{and} \quad \lambda(Q) - \lambda(R) = \int_z^0 -\frac{gF}{y_L - F} dx.$$

Therefore,

$$\lambda(Q) - \lambda(P) = \int_0^z \left\{ \left(-\frac{gF}{y_U - F} \right) - \left(-\frac{gF}{y_L - F} \right) \right\} dx$$

or

$$\Delta(z) = \frac{1}{2}[y^2(Q) - y^2(P)] = \int_0^z F(x) \frac{d}{dx} (y_U - y_L) dx. \quad (5)$$

Notice that if for some value of z , $\Delta(z) = 0$, then the orbit passing through $(z, F(z))$ is periodic, for then $y(Q) = -y(P)$, and by the uniqueness, and mirror symmetry of trajectories through the origin, (when F and g are odd) the path will be closed.

Thus the proof of the existence of an infinite number of limit cycles now depends on demonstrating that $\Delta(z)$ has an infinite number of zeros.

Setting $d/dx(y_L - y_U) = h(x, z)$ we have $\Delta(z) = -\int_0^z F(x) h(x, z) dx$. Observe that $h(x, z) \geq 0$ for $0 \leq x \leq z$.

We need the result of D'heedene and Comstock (which we call Lemma I):

If $h(x, z)$, for fixed z , is a nondecreasing function of x (for $0 \leq x \leq z$) then

$$\text{sign } \Delta(z_n) = (-1)^n.$$

The crucial part of the argument now lies in showing that $h(x, z)$ is monotone increasing in x , for then, by the continuity of $\Delta(z)$, a periodic orbit exists between successive zeros of $F(x)$.

THEOREM 1. *If H_1 is satisfied, and $g(-x) = g(x)$ with $g \in C^1(-\infty, \infty)$ and $4g'(x) \geq f^2(x)$ for all x , then (3) has an infinite number of limit cycles.*

Proof.

$$h(x, z) = \frac{d}{dx} y_L - \frac{d}{dx} y_U.$$

Now it is sufficient for

$$h' = y_L'' - y_U'' \geq 0$$

to have $y_L'' \geq 0$ and $y_U'' \leq 0$ separately.

Consider first

$$y_U'' = \frac{d}{dx} \left(-\frac{g}{y_U - F} \right) = - \left\{ \frac{g'(x)\xi^2 + g(x)f(x)\xi + g^2(x)}{\xi^3} \right\},$$

where $\xi = y_U - F(x) \geq 0$ in $0 \leq x \leq z$.

Since $g^2(x)f^2(x) \leq 4g'(x)g^2(x)$ by hypothesis, it follows that the discriminant (of the quadratic in ξ) is negative, and therefore $y_U'' \leq 0$.

A similar argument shows that $y_L'' > 0$.

Remark. Notice that for (1) or (2), the condition $4g'(x) \geq f^2(x)$ becomes, $\mu^2 \leq 4$. In general, if a parameter μ is introduced, this condition may provide an upper bound on μ .

Unfortunately, the trajectories behave quite differently for large and for small μ ; for example, $y_U'' \leq 0$ is not generally satisfied for μ large.

It was pointed out, however, by D'heedene and Comstock that $h'(x, z) \geq 0$ still holds for trajectories sufficiently remote from the origin.

THEOREM 2. *If H_1 and H_2 are satisfied, then (3) has an infinite number of limit cycles.*

Proof. It is sufficient to show that $h'(x, z) \geq 0$ for $z \geq x^*$, for then, by Lemma I, the theorem follows.

Now

$$h' = y_L'' - y_U'', \text{ and } y_U' = -\frac{g}{y_U - F} = -g'\xi, \quad y_L' = -\frac{g}{y_L - F} = \frac{g}{\eta},$$

where $\xi = y_U - F \geq 0$ and $\eta = F - y_L \geq 0$.

Performing the differentiations, we obtain

$$h' = \frac{\xi + \eta}{\xi^3 \eta^3} [g^2(\eta - \xi)^2 + g^2 \xi \eta + g' \xi^2 \eta^2 + g f \xi \eta (\eta - \xi)].$$

Now h' will certainly be nonnegative if

$$\psi = g^2(x) + g'(x) \xi \eta + g(x) f(x) (\eta - \xi)$$

is nonnegative.

One has from (4), that

$$|\eta - \xi| = |y_U + y_L - 2F| \leq 4A.$$

Also

$$\int_0^x g(u) g'(u) du \geq a \int_0^x g(u) du = aG(x).$$

Therefore,

$$g^2(x) \geq 2aG(x). \quad (6)$$

On using $|f(x)| \leq B$, one gets

$$\psi \geq 2aG(x) + a\xi\eta - 4ABg(x). \quad (7)$$

We now require a lower bound on $\xi\eta$.

If x lies in R_1 , the region where

$$[2\{G(z) - G(x)\}]^{1/2} \geq 2A$$

then (4) provides a positive lower bound on ξ and η , viz.,

$$\xi \geq [2\{G(z) - G(x)\}]^{1/2} - 2A,$$

and likewise for η . Therefore,

$$\xi\eta \geq 2[G(z) - G(x)] - 4A[2\{G(z) - G(x)\}]^{1/2} + 4A^2$$

or

$$\xi\eta \geq 2G(z) - 2G(x) - 4A[2G(z)]^{1/2} + 4A^2.$$

Using this in (7) yields

$$\psi \geq 2aG(z) - 4aA[2G(z)]^{1/2} - 4ABg(x) + 4aA^2.$$

Using $g(x) \leq g(z)$ and (6) in the form $g(z)/a^{1/2} \geq [2G(z)]^{1/2}$ results in

$$\psi \geq 2ag(z) \left[\frac{G(z)}{g(z)} - \frac{2A}{a}(B + \sqrt{a}) \right] + 4aA^2$$

or

$$\psi > 0 \quad \text{for } z > x^*, \text{ by } H_2(b).$$

If, however, x lies in R_2 , the region where

$$\{2[G(z) - G(x)]\}^{1/2} < 2A,$$

the lower bound obtained above, for ξ and η , is negative. In this region, we use

$$G(x) > G(z) - 2A^2, \quad g(x) \leq g(z)$$

and $\xi\eta \geq 0$ to obtain, from (7),

$$\psi \geq 2a[G(z) - 2A^2] - 4ABg(z),$$

putting $\xi\eta = 0$ (since $\xi\eta \geq 0$).

On invoking H_2 , viz.,

$$G(z) \geq \frac{2A}{a}(B + a^{1/2})g(z) \quad \text{for } z > x^*$$

yields

$$\psi \geq 4Aa^{1/2}g(z) - 4aA^2 = 4Aa^{1/2}[g(z) - a^{1/2}A].$$

Since $g(z) = \int_0^z g'(x) dx \geq az$, then

$$g(z) - a^{1/2}A \geq az - a^{1/2}A > 0,$$

whenever $z > A/a^{1/2}$, which is certainly satisfied for $z > x^*$.

Consequently, for orbits so large that $z > x^*$, then $h'(x, z) \geq 0$, whether x is in R_1 or in R_2 , and $\text{sign } \Delta(z_n) = (-1)^n$, by Lemma I for $z > x^*$. Hence an infinite number of limit cycles exist

3. PERIODIC ORBITS FOR $F(x)$ EVEN, $g(x)$ ODD

Let

h_1 :

- (a) $F(x)$ satisfy a Lipschitz condition for every finite x interval,
- (b) $F(x) = F(-x)$,
- (c) $F(x) < 0$ for $x > 0$.

and h_2 :

- (a) $g(x)$ satisfy a Lipschitz condition for every finite x interval,
- (b) $xg(x) > 0$ for $x \neq 0$,
- (c) $g(x) = -g(-x)$,
- (d) $G(\pm\infty) = \infty$.

THEOREM 3. *The system (3) possesses a nondenumerable set of periodic solutions, if the hypotheses h_1 and h_2 are satisfied.*

Proof. We first show that any solution trajectory of (3) starting on the negative y axis will intersect the y axis again. Next, we prove that the orbits described by (3) are symmetrical about the y axis, closed, hence periodic.

A solution starting at $(0, -y_0)$, $y_0 > 0$, will cross the closed contours $\lambda(x, y, 0) = C$ toward their interiors since $\dot{\lambda} = -g(x)F(x) \leq 0$ for $x \leq 0$.

Next, observe that the motion in reverse time $\tau = -t$, with $X = -x$, is governed by $X' = y - F(X)$, $y' = -g(X)(') \equiv d/d\tau$, the same equations as (3). Thus the orbits described by (3) have mirror symmetry about the y axis. A solution beginning at $(0, -y_0)$ will intercept the y axis again at $(0, y_1)$ and return to $(0, -y_0)$, resulting in a closed, hence periodic orbit. It is clear that a periodic orbit passes through every point on the negative y axis. There are an uncountable number of limit cycles. Furthermore, it follows from the uniqueness of the solutions that the negative y axis is mapped by the orbits into a nondenumerable set of points on the positive y axis ($y_1 > 0$).

COROLLARY 3.1. *Let $h_1(c)$ be replaced by $h_1(c')$: $F(x) > 0$ for $x > 0$, all other hypotheses of Theorem 3 still holding, then Theorem 3 is still true. A periodic orbit will pass through each point on the positive y axis.*

COROLLARY 3.2. *Let $h_2(d)$ be replaced by $h_2(d')$: $G(\pm\infty) = Y_0^2$, $Y_0 > 0$, all other hypotheses of Theorem 3 still holding, then Theorem 3 is still true. A periodic solution will pass through each point in the interval $(0, -Y_0)$ on the negative y axis.*

AN EXAMPLE. The system

$$\dot{x} = y + x^{2n}, \quad \dot{y} = -x^{2n-1} \quad (n \text{ a positive integer}) \quad (8)$$

satisfies all the conditions of Theorem 3. The corresponding phase equation can be integrated exactly; the solution trajectories are given by

$$x^{2n} = C \exp(-2ny) + \frac{1 - 2ny}{2n}. \quad (9)$$

If $-1/2n < C < 0$, the solutions are periodic, if $C \geq 0$ aperiodic. Periodic orbits intercept the negative y axis in $(-\infty, 0)$ and the positive y axis in $(0, 1/2n)$, whereas aperiodic orbits intercept the y axis once, in $(1/2n, +\infty)$.

THEOREM 4. *Let $h_1(c)$ be replaced by $h_1(c'')$: $|F(x)| \leq A$ for all x , all other hypotheses of Theorem 3 still holding, then the periodic orbits cover the $x - y$ plane.*

Proof. Starting with any initial point (x_0, y_0) in the plane the trajectory through that point is bounded by the oval

$$\lambda(x, y, -A) = \frac{(y + A)^2}{2} + G(x) = C \quad \text{for } x \geq 0$$

since $\dot{\lambda} = -g(x)[F(x) + A] \leq 0$ for $x \geq 0$, and by the closed nested ovals $\lambda(x, y, A) = C$ for $x \leq 0$, since $\dot{\lambda} = -g(x)[F - A] \leq 0$ for $x \leq 0$. The families of ovals $\lambda(x, y, -A) = C$, $\lambda(x, y, A) = C$ thus guide solutions to the axes.

It follows, from the mirror symmetry of orbits about the y axis, that a solution starting at $(x_0, 0)$, with $x_0 > 0$, will intercept the negative y axis at $(-x_0, 0)$, and return to $(x_0, 0)$, resulting in a closed, hence periodic, orbit. By proper choice of x_0 , a periodic orbit can be made to pass through any point in the plane. Q.E.D.

COROLLARY 4.1. *Let $F(x)$ be oscillatory and let the hypotheses of Theorem 4 be satisfied. Then the periodic orbits cover the $x - y$ plane.*

REFERENCES

1. H. HOCHSTADT AND B. H. STEPHAN, On the limit cycles of $\ddot{x} + \mu \sin \dot{x} + x = 0$, *Arch. Rational Mech. Anal.* **23** (1967), 369-379.
2. R. N. D'HEEDENE, "For all Real μ , $\ddot{x} + \mu \sin \dot{x} + x = 0$ has an Infinite Number of Limit Cycles," *J. Differential Equations* **5** (1969), 564-571.
3. C. COMSTOCK, On the limit cycles of $y'' + \mu F(y') + y = 0$, *J. Differential Equations* **8** (1970), 173-179.